



LIQUID IRON

For Correction or Prevention of Iron Deficiency in Most Crops

GUARANTEED ANALYSIS

Sulfur (S)......4.00%

4.00% Combined Sulfur (S)

Iron (Fe)..... 5.00%

5.00% Chelated Iron (Fe)

5.00% Water Soluble Iron (Fe)

Derived from: Iron(II) Sulfate and Iron Lignin Sulfonate

<u>Net Weight</u>: Jug: 25lbs (11.34 kg) Cs.2x2.5: 51lbs (22.68 kg) <u>Net Contents</u>: Jug: 2.5 gal (9.46 liters) Cs. 2x2.5: 5 gal (18.92 liters)

WHAT TO DO IN CASE OF CONTACT

If Ingested: Give large amounts of water or milk, if conscious. Contact a physician immediately. Eye Contact: Flush eyes with clean water for 15 minutes. Seek medical attention immediately.

Skin Contact: Remove contaminated clothing and wash skin with soap and water. Seek medical attention if irritation develops or persists.

If Inhaled: Move to fresh air and seek medical attention if irritation develops or persists.

For additional information call CHEMTREC 800-424-9300

If there is contact made with spray solution containing pesticides and/or spray adjuvants, follow the "Statement of Practical Treatment" on the product labeling.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

READ LABEL - FOLLOW DIRECTIONS!

Refer to MSDS for health, safety and environmental information. Keep out of Reach of Children

GENERAL INFORMATION

SUPER-KEY LIQUID IRON is a proprietary blend of soluble micronutrient(s) and an organic complexing agent so as to enhance and maintain availability and accelerate uptake by crops.

SUPER-KEY LIQUID IRON is non-phytotoxic and can be applied foliar to most crops. It is recommended for use throughout the season in a program of multiple applications for greater efficiency. This provides a continuous supply of vital nutrients at or near the time they are needed most.

SUPER-KEY LIQUID IRON is environmentally safe, made with only "clean" raw materials free from heavy metal contamination.

DIRECTIONS FOR USE

SUPER-KEY LIQUID IRON can be added to most nitrogen solutions or fluid fertilizers and impregnated on dry blend fertilizer. It can be tank mixed with most pesticides. Always make a small-scale test to assure compatibility. It can be soil or foliar applied, broadcast or banded over the row and injected through irrigation systems. CAUTION: DO NOT USE IN HIGH PHOSPHATE "STARTER" OR "POP-UP" FERTILIZER MATERIALS SUCH AS 10-34-0.

SUPER-KEY LIQUID IRON will disperse in water with minimal agitation. It serves as a buffer when used in tank mixes with pesticides. Follow the mixing sequence: 1. Water; 2. SUPER-KEY LIQUID IRON; 3. Pesticide.

RECOMMENDED RATES

TURF APPLICATION			
Application		Rate Per 1,000 sq. ft.	Frequency
Tees, Greens, All Types of		3 – 4 oz. (90-120 ml)	Apply every 14-21 days
Warm and Cool Season Grasses			
Fairways		3 – 4 oz. (90-120 ml)	Apply every 30 days
TREES, SHRUBS & ORNAMENTALS			
Soil Injection:	Apply 4 oz. per inch of trunk DBH in 2 gallons of water or as needed		
	according to soil and tissue deficiencies.		
Foliar Spray:	20 oz in 100 gallons of water (4-5 L in 400 L water)		
Soil Drench	3 ounces per inch of trunk DBH in 2 gallons of water. Thoroughly		
Corrective:	drench soil around area (90 ml per 8 L water)		
Soil Drench	2 oz per inch of trunk DBH in 2 gallons of water (60 ml per 8 L water)		
Maintenance:			
Budding Plants	Mix Super-Key Liquid Iron at a rate of 3 o in 4 gallons of water and		
& Perennials:	spray foliage to run off. Apply as needed throughout the growing		
	season.		

The optimum rate and timing of application will depend on local conditions and should always be made as a result of soil and/or plant tissue analysis in consultation with your local agricultural authorities. In the absence of test results, use the following as a guide.

This product may be corrosive to aluminum, mild steel and brass. Store in HDPE, fiberglass or stainless steel containers. Use only stainless steel, P.V.C. or polypropylene fittings.

Storage: Store in a cool place in original container in a locked storage area inaccessible to children, pets, domestic animals or wildlife. Container Disposal: Do not reuse empty containers. Triple rinse (or equivalent) then offer for recycling or disposal by procedures approved by state or local authorities.

CONDITIONS OF SALE: The Doggett Corp. (Company) warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal use. This warranty does not extend to the use of this product contrary to label instruction, or under abnormal use conditions, or under conditions not reasonably foreseen by the company. THE COMPANY DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. THE COMPANY SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND THE COMPANY'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND SELLER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. THE COMPANY DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

The manufacturer disclaims all responsibility for damage to plants and equipment through the use of this product whether used in accordance with directions or not.